

IFUP-TH 68/99

LPT-ENS-99/59

ADM approach to 2+1 dimensional gravity ¹

Pietro Menotti ²

*Dipartimento di Fisica dell'Università, Pisa 56100, Italy and
INFN, Sezione di Pisa*

Domenico Seminara³

*Laboratoire de Physique Théorique, École Normale Supérieure⁴,
F-75231, Paris CEDEX 05, France*

December 1999

¹Contribution to Third Meeting on Constrained Dynamics and Quantum Gravity, Villasimius, September 13-17, 1999

²This work is supported in part by M.U.R.S.T.

³CEE Post-doctoral Fellow under contract FMRX CT96-0045

⁴Unité Mixte associée au Centre de la Recherche Scientifique et à l'École Normale Supérieure.

Abstract

The canonical ADM equations are solved in terms of the conformal factor in the instantaneous York gauge. A simple derivation is given for the solution of the two body problem. A geometrical characterization is given for the apparent singularities occurring in the $\mathcal{N}$ -body problem and it is shown how the Garnier hamiltonian system arises in the ADM treatment by considering the time development of the conformal factor at the locations where the extrinsic curvature tensor vanishes. The equations of motion for the position of the particles and of the apparent singularities and also the time dependence of the linear residues at such singularities are given by the transformation induced by an energy momentum tensor of a conformal Liouville theory. Such an equation encodes completely the dynamics of the system.

1 Introduction

The problem of providing solutions to $2 + 1$ dimensional gravity [1] has been approached along different lines. In absence of particles the dynamics becomes non trivial only for closed universes of genus higher or equal to one. Moncrief [2] and Hosoya and Nakao [3] gave the hamiltonian treatment of this problem providing the complete reduction of the hamiltonian to the physical parameters i.e. the moduli of the time slices. This procedure was obtained in the York $K = t$ gauge; in the case of the torus the problem can be dealt with explicitly while for higher genus, even if the problem is well defined we do not possess an explicit form for the reduced hamiltonian.

A different approach was put forward by 't Hooft [4] by describing the evolving Cauchy surfaces in terms of polygonal tiles which join along segments where the extrinsic curvature is singular. The dynamics of the system is codified in transition rules which intervene when e.g. the length of a side of a polygon goes to zero or when a particle collides with a side of a polygon. Such an approach is applicable both in presence and in absence of particle and it is suitable for an algorithmic or numeric approach. A similar but different approach was given by Waelbroek [5].

In presence of particles progress was made in the papers by Bellini, Ciafaloni and Valentoli [6] and by Welling [7] in the first order formalism by going over to the instantaneous ($K = 0$) gauge; in these papers it was shown that the problem is equivalent to a Riemann-Hilbert problem. It can be solved explicitly for the case of two particles, in terms of hypergeometric functions. For three or more particles one encounters a feature known in the mathematical literature as apparent singularities.

In this paper we shall consider $2 + 1$ dimensional gravity in presence of massive particles by exploiting the hamiltonian formulation. Thus our setting is a second order formalism and the basic equations are those derived systematically from the variation of the ADM action. The reason for such a development is to give a treatment which resides completely within the canonical framework. All results obtained in [6] [7] come out in simple fashion from such an approach; moreover we shall prove that the Garnier hamiltonian system for the apparent singularities is a direct outcome of the canonical ADM formalism. Again the exploitation of the instantaneous York gauge plays a major role; the technical advantage of such a gauge is to reduce an equation of sinh-Gordon to one of Liouville type to which powerful methods of complex analysis apply. Unfortunately the applicability of such a gauge is restricted to open universes.

We shall see that the described approach provides an elementary way to solve the two body problem: the exact solution of the motion can be derived even before the explicit computation of the metric; combined with the qualitative knowledge of the asymptotic metric it gives a complete description of the scattering. The exact metric is obtained by solving a Liouville equation, leading of course to the same results obtained in [6] and [7].

One of the aims of presenting a canonical solution of the problem is to provide a framework for the quantization of the theory even if here we shall confine ourselves to the classical problem.

Quantization schemes for $2 + 1$ dimensional gravity in absence of particles have been proposed in [8, 9, 10] and in presence of particles in [11, 12, 13].

The key quantity of our approach will be the conformal factor which describes the metric of the time slices. The lapse function N is given by the derivative of such conformal factor

with respect to the total energy and the shift functions are given in terms of derivatives of the lapse function.

As mentioned above, with more than two particles there are apparent singularities and one has to provide evolution equations for their positions and residues. We show that all this information is contained in the ADM equation for the time derivative of the conformal factor. In fact by exploiting Schwarz's relation between the coefficient of the fuchsian differential equation and the reduced conformal factor one obtains the Garnier equation by equating the residues at the polar singularities at the position of the apparent singularities.

In the present paper we shall outline the main features of the approach; technical details are found in ref.[14].

2 The action

The action for the gravitational field including the boundary terms is given by [15]

$$\begin{aligned} \frac{16\pi G_N}{c^3} S_{Grav} = & \int_{\mathcal{M}} dt d^2x \sqrt{\mathbf{g}} \mathcal{R} + \\ & 2 \int_{\Sigma_0}^{\Sigma_1} d^2x \sqrt{\mathbf{g}} K + 2 \int_B d^2x \sqrt{-\gamma} \Theta + \\ & 2 \int_{B_0}^{B_1} dx \sqrt{\sigma} \eta \end{aligned} \quad (1)$$

where $\mathbf{g}_{\mu\nu}$, $\mathcal{R}$ are the $2+1$ dimensional metric and curvature, g_{ij} the 2-dimensional metric of the constant time slices Σ_t ; Σ_0 and Σ_1 the initial and final time slices; K_{ij} the second fundamental form of the time slices Σ_t and Θ_{ij} the second fundamental form of the lateral boundary B whose volume form is $\sqrt{-\gamma}$; n^μ is the future pointing unit normal to the time slices and u^μ the outward pointing unit normal to B ; $B_t = \Sigma_t \cap B$, $\sqrt{\sigma}$ is the volume form induced on B_t and $\sinh \eta = n_\mu u^\mu$. The boundary terms are going to play a fundamental role in the computation of the reduced action; this paper however will be based completely on the equations of motion which do not depend on the explicit form of the boundary terms.

In presence of particles we must add to (1) the term

$$- \int dt \sum_n m_n \sqrt{-\mathbf{g}_{\mu\nu}(q_n) \dot{q}_n^\mu \dot{q}_n^\nu} \quad (2)$$

with $m_n = \frac{16\pi G_N m_n}{c^2}$, $0 < m_n < 4\pi$. In the ADM notation where the metric is written as [16]

$$ds^2 = -N^2 dt^2 + g_{ij}(dx^i + N^i dt)(dx^j + N^j dt) \quad (3)$$

the volume term apart for a divergence, can be rewritten in terms of the canonical variables as [16, 15, 17]

$$\int_{\mathcal{M}} dt d^2x \left[\pi^{ij} \dot{g}_{ij} - N^i H_i - NH \right] \quad (4)$$

where we have introduced the momenta π^{ij} canonically conjugate to g_{ij} . Similarly the matter action takes the form

$$\int dt \sum_n \left(P_{ni} \dot{q}_n^i + N^i(q_n) P_{ni} - N(q_n) \sqrt{P_{ni} P_{nj} g^{ij}(q_n) + m_n^2} \right). \quad (5)$$

3 The constraints

In (2+1)–dimensions, the great advantage of a York-type gauge, $\partial_i K = 0$, is to ensure, when combined with the conformal gauge $g_{ij} = \exp(2\sigma)\delta_{ij}$, the decoupling of the traceless part π^{Tij} of the momentum from the spatial metric in the diffeomorphism constraints⁵. Indeed, the diffeomorphism constraints,

$$H_i = -2\sqrt{g}\nabla_j \frac{\pi^j_i}{\sqrt{g}} - \sum_n \delta^2(x - q_n) P_{ni} = 0, \quad (6)$$

become

$$\partial_z \pi^{Tz}_{\bar{z}} = -\frac{1}{2} \sum_n \delta^2(x - q_n) P_{nz}$$

and

$$\partial_{\bar{z}} \pi^{T\bar{z}}_z = -\frac{1}{2} \sum_n \delta^2(x - q_n) P_{n\bar{z}},$$

which are solved by

$$\pi^{T\bar{z}}_z = -\frac{1}{2\pi} \sum_n \frac{P_{nz}}{z - z_n} + P_{\bar{z}}(z) \quad (7)$$

⁵This decoupling reflects the fact that the divergence of a traceless symmetric tensorial density of rank two is Weyl-invariant in two dimensions.

and its complex conjugate expression. Here we introduced the complex coordinates z and $\bar{z}$ with $z = x + iy$ and $P_{\bar{z}}^{\bar{z}}(z)$ is an *entire* function. In the following, we shall adopt the *maximally slicing* gauge $K = 0$, where π^{ij} coincides with its traceless part so that so that eq.(7) holds for $\pi_{\bar{z}}^{\bar{z}}$ and its complex conjugate for π_z^z .

We come now to the Hamiltonian constraint,

$$H = \frac{1}{\sqrt{g}} \left[\pi^a_b \pi^b_a - (\pi^c_c)^2 \right] - \sqrt{g} R \quad (8)$$

$$+ \sum_n \delta^2(z - z_n) \sqrt{m_n^2 + P_{ni} g^{ij} P_{nj}} = 0.$$

If we integrate the l.h.s of eq.(8) over a slice of constant time, an elementary application of the Gauss-Bonnet theorem shows that the condition $K = 0$ is consistent only with open universes or with closed universes of genus 0. In the latter case consistency can be achieved only for static universes as it will be shown in Sect.4. Moreover, for open universes, eq.(8) determines the asymptotic behavior of the momenta π^i_j in terms of that of σ . In fact if we assume that the asymptotic metric has to be conical, $e^{2\sigma} \sim (z\bar{z})^{-\mu}$, $\mu > 0$, the above equation implies that

$$\int d^2z \, 2\pi_z^z \pi_{\bar{z}}^{\bar{z}} \exp(-2\sigma) < \infty.$$

This, in turn, entails that $\pi_{\bar{z}}^{\bar{z}}$ at infinity vanishes faster than $1/z$. Thus, the requirement of having a conical metric at infinity sets the function $P_{\bar{z}}^{\bar{z}}(z)$ to zero and in addition it imposes that $\sum_n P_{nz} = 0$. Because of the explicit form of the solution (7), we actually end up with a $\pi_{\bar{z}}^{\bar{z}}$ that goes to zero as $1/z^2$. As a consequence, if we define, mimicking the four dimensional counterpart, the total momentum P_{tot} of the system as the surface integral of π^i_j , we are led to conclude that P_{tot} vanishes identically. It is worth noticing that in the present gauge P_{tot} is simply equal to $\sum_n P_n$ and no gravitational correction appears. The clash in $(2 + 1)$ -dimensions between the boundary conditions ensuring finite energy and those ensuring finite momentum was first noticed in [18, 19, 20]. In particular, in [19] the origin of this conflict was traced back to the impossibility of performing asymptotic boosts without introducing spurious singularities in the metric.

We return now to eq.(8). It is not difficult to show [14] that $e^{-2\sigma}$ vanishes on the particle singularities and as a result the hamiltonian constraint assumes the form, after

setting $2\tilde{\sigma} = 2\sigma - \ln(2\pi_{\bar{z}}^{\bar{z}}\pi_z^z)$

$$2\Delta\tilde{\sigma} = \tag{9}$$

$$-e^{-2\tilde{\sigma}} - \sum_B 4\pi\delta^2(z - z_B) + \sum_n \delta^2(z - z_n)m_n.$$

The above equation is an inhomogeneous Liouville equation. The sources are given by the particles and by some other auxiliary sources whose position z_B are the zeros of $\pi_{\bar{z}}^{\bar{z}}$ and their strength is fixed to -4π . The conformal factor σ is actually regular at the points z_B and for this reason we shall refer to them as *apparent* singularities.

We remark that the reduction of the hamiltonian constraint to a Liouville equation is the result of the $K = 0$ condition. The $K = \text{const} \neq 0$ gauge gives rise, as seen from eq.(8) to an equation of the sinh-Gordon family which is of much more complex nature.

Finally we note that the conformal gauge i.e. $g_{ij} = e^{2\sigma}\delta_{ij}$ is still not a complete gauge fixing. One is allowed to perform analytic transformation on z which leave, in the case of open universe, the point at infinity fixed i.e. time dependent translations, rotations and dilatations. This freedom can be exploited e.g. to keep one particle fixed at $z_1 = 0$ and an other fixed at $z_2 = 1$. In general such a fixing is equivalent to a frame which rotates at infinity.

4 Equations for the lapse and shift functions

Before discussing the solutions of the inhomogeneous Liouville equation we shall write down the equations for N and N^i ; these are an outcome of requiring that the gauge conditions are preserved in the Hamiltonian evolution. Combining the canonical equation for $\dot{\pi}^{ij}$ and the one for $\dot{g}_{ij}$ the relation $K = 0$ (or equivalently $\pi^i_i = 0$) takes the form

$$\Delta N = e^{-2\tilde{\sigma}} N. \tag{10}$$

Again we have exploited the vanishing of $e^{-2\sigma}$ at the particle sources [14]. Since we have chosen the conformal gauge, the traceless part of the equation for $\dot{g}_{ij}$ determines N^i , i.e.

$$\partial_{\bar{z}} N^z = -\pi_{\bar{z}}^z e^{-2\sigma} N \tag{11}$$

and its complex conjugate. These complete the list of our equations for the metric. By applying Gauss theorem to eq.(10) one easily derives the asymptotic behavior of N for large $|z|$

$$N \sim \ln(z\bar{z}). \quad (12)$$

Such an asymptotic behavior of the lapse function is typical of the instantaneous York gauge and should be contrasted with the behavior $N \rightarrow \text{const.}$ of the familiar DJH gauge [1]. The behavior (12) holds both when the plane describes the time slice of an open universe and when it describes the sphere; but in the latter case in order for the scalar N to be well defined on the Riemann sphere it should go to a constant at infinity. This occurs only for $\pi^i_j \equiv 0$ i.e. when the equation for N reduces to $\Delta N = 0$. Thus we see that except for the trivial stationary case in which $P_n = 0$ the York instantaneous gauge applies only to open universes.

5 The Liouville equation

As is well known, the problem of solving the inhomogeneous Liouville equation (9) can be reduced to that of solving a fuchsian second order differential equation of projectively canonical type [21]

$$y''(\zeta) + Q(\zeta)y(\zeta) = 0 \quad (13)$$

where

$$Q(z) = \sum_n \left[\frac{1 - \mu_n^2}{4(z - z_n)^2} + \frac{\beta_n}{2(z - z_n)} \right] + \sum_B \left[-\frac{3}{4(z - z_B)^2} + \frac{\beta_B}{2(z - z_B)} \right]. \quad (14)$$

If we define the function $f(z)$ as the ratio of two independent solutions of eq. (13),

$$f(z) = \frac{ky_1(z)}{y_2(z)}, \quad (15)$$

the solution of eq. (9) is built from the Poincaré metric for the pseudosphere by performing on it the conformal diffeomorphism defined by eq. (15)

$$e^{-2\tilde{\sigma}} = \frac{8f'(z)\bar{f}'(\bar{z})}{(1-f(z)\bar{f}(\bar{z}))^2}. \quad (16)$$

The poles in (14) include both particle and apparent singularities. The residues of the second order poles in $Q(z)$ are directly related to the strength of the delta singularities in eq.(9): in fact, $\mu_n = m_n/4\pi$ while the second order residue at the apparent singularities is fixed to be $-3/4$. The latter value is required by the regularity of 2σ at z_B . The determination of the first order residues is more difficult and it is related to the knowledge of the explicit solution of the Riemann-Hilbert problem associated to (13). They are fixed in principle by the requirement that the conformal factor σ be monodromic in the complex plane. As eq.(16) can be rewritten in the form

$$e^{-2\tilde{\sigma}} = \frac{8w_{12}\bar{w}_{12}}{(y_2\bar{y}_2 - k\bar{k} y_1\bar{y}_1)^2} \quad (17)$$

where w_{12} is the constant wronskian, it is seen that such a requirement is equivalent to the imposition that the vector (ky_1, y_2) transforms under a representation of $SU(1,1)$ when one encircles a singularity in the complex plane. It provides the relation between the geometric structure approach [22] and the one here described.

An important feature in the solution of eq.(9) is that one has to give the residue of the singularity at infinity and this information is not contained in the parameters appearing in the sources. In different words, in solving eq.(9) the total energy of the system has to be specified as an arbitrary parameter M because the positions and the momenta of the particles at the initial time are not enough to determine its value. The above conclusion might seem counterintuitive especially if we compare it with what happens in the DJH *geometrical* approach, where the knowledge of all z_i and p_i fixes M completely through a nonlinear composition rule for the momenta. This apparent mismatch simply reflects their choice to have, in contrast with the present approach [6, 7, 14], a *flat* (not globally defined) metric. As a consequence, in their approach, all the dynamical information is shifted into the particle variables. In our approach a complete set of initial conditions for the conformal

factor $\tilde{\sigma}$ is instead provided by any unconstrained parameterization⁶ of the rational function $Q(z)$: an example will be given in sec. 9.

The conformal factor $2\tilde{\sigma}$ plays the key role in all subsequent developments; once the solution of eq.(9) is known the equations for N and N^i are solved as follows. As the sources in eq.(9) do not depend on the total energy M , by taking the derivative with respect to M we find a solution to the homogeneous equation (10) for N i.e. apart for an arbitrary normalization constant we have

$$N = \frac{\partial(-2\tilde{\sigma})}{\partial M}. \quad (18)$$

As for the equation (11) multiplying it by $2\pi_{\bar{z}}^z$ and using eq.(10) and taking into account that $\pi_{\bar{z}}^z$ is a function of z we get

$$N^z = -\frac{2}{\pi_{\bar{z}}^z(z)}\partial_z N + g(z) \quad (19)$$

where $g(z)$ is a function of z . A similar solution holds for $N^{\bar{z}}$. The function $g(z)$ has to be chosen as to cancel the poles generated by the zeros of $\pi_{\bar{z}}^z(z)$, which occur only in presence of three or more particles, and if we are interested in describing a reference system which does not rotate at infinity $g(z)$ has to be chosen as to give $N^z < |z|$ at infinity. We shall see in sect.7 that, due to $\sum_n P_{nz} = 0$, such a function can diverge at infinity at most linearly. As we shall see $g(z)$ encodes the time evolution of all the quantities of the problem.

6 The particle equations of motion

The canonical equations for the particle positions z_n and momenta P_n are obtained by varying the action with respect to z_n and P_n and are given by

$$\dot{z}_n = -N^i(z_n, t) = -g(z_n), \quad (20)$$

⁶ In fact the β_n and β_A are not all independent: In the next section we shall see that fuchsianity and the value of the residue at infinity impose two sum rule on them. Moreover, the requirement that conformal factor σ be regular at z_A fixes the β_A in terms of the other parameters by means of the so-called *no-log* condition, see sec.9.

$$\dot{P}_{nz} = P_{na} \frac{\partial N^a}{\partial z} - m_n \frac{\partial N}{\partial z} \quad (21)$$

and their complex conjugate expressions. We note that $\dot{z}_n$ is simply given in terms of the meromorphic function $g(z)$ as in eq.(19) $\pi^{\bar{z}}_z$ diverges at the particle positions while $\partial_z N$ stays finite. It is of interest that the r.h.s of eq.(21) depends only on the function $g(z)$ and on the linear residue β_n at the particle singularity. In fact by using the behavior of the solutions eq.(13) at the particle singularities, one obtains

$$\dot{P}_{nz} = 4\pi \frac{\partial \beta_n}{\partial M} + P_{nz} g'(z_n). \quad (22)$$

The fact that the time evolution is governed by the function $g(z)$ will be the common denominator of the following developments.

7 Conservation laws

For a fuchsian differential equation of type (13) the following Fuchs relations hold [23]

$$\begin{aligned} \sum_n \beta_n + \sum_B \beta_B &= 0; \quad 1 - \mu_\infty^2 = \\ &= \sum_n (1 - \mu_n^2 + 2\beta_n z_n) + \sum_B (-3 + 2\beta_B z_B). \end{aligned} \quad (23)$$

The former expresses the fact that the function $Q(z)$ behaves as $1/z^2$ at infinity (*Fuchs condition*); the latter provides the second order residue of $Q(z)$ at infinity (*the total mass* of the system) in terms of the other parameters.

Below, we shall show that these conditions imply some non-trivial conservation laws. We start recalling the equation of motion for $\pi^{\bar{z}}_z$

$$\dot{\pi}^{\bar{z}}_z = 2e^{2\sigma} \partial_z (e^{-2\sigma} \partial_z N) + 2\pi^{\bar{z}}_z \partial_z N^z + N^z \partial_z \pi^{\bar{z}}_z \quad (24)$$

which by means of the eqs.(18) and (19) can be recasted as follows

$$\dot{\pi}^{\bar{z}}_z = -4\partial_M Q(z) + \partial_z \pi^{\bar{z}}_z g + 2\pi^{\bar{z}}_z \partial_z g. \quad (25)$$

We have used a standard result about the Liouville equation that permits to connect $\tilde{\sigma}$ to $Q(z)$

$$Q(z) = -\frac{1}{2}[\partial_z^2(2\tilde{\sigma}) + \frac{1}{2}(\partial_z(2\tilde{\sigma}))^2]. \quad (26)$$

In a field theoretical language $Q(z)$ is proportional to the energy momentum tensor associated to the Liouville equation.

If we integrate both sides of eq.(25) over a closed path in the complex plane that encircles all the singularities, we find

$$\begin{aligned} \frac{d}{dt} \sum_n P_{nz} &= 4\pi \partial_M (\sum_n \beta_n + \sum_B \beta_B) - \\ &2\pi \oint dz (\partial_z \pi_{\bar{z}} g + 2\pi_{\bar{z}} \partial_z g). \end{aligned} \quad (27)$$

Since both the sum over momenta and that over the first order residues vanish, the remaining integral must vanish as well. Recalling that $\pi_{\bar{z}} \simeq 1/z^2$ for large radius, we must conclude that g can grow at most linearly at infinity. Moreover, if we combine this information with the fact that the only singularities of g at finite radius can be poles, we find that g must be a rational function.

Next, we multiply eq.(25) by z and we integrate again over a closed path in the complex plane to obtain

$$\begin{aligned} \frac{d}{dt} \sum_n z_n P_{nz} &= \\ 4\pi \partial_M (\sum_n z_n \beta_n + \sum_B z_B \beta_B) &= 1 - \frac{M}{4\pi}. \end{aligned} \quad (28)$$

The previous equation combined with its complex conjugate gives both the law of conservation of angular momentum

$$\dot{L} = \frac{d}{dt} \sum_n i(z_n P_{nz} - \bar{z}_n P_{n\bar{z}}) = 0 \quad (29)$$

and the “generalized conservation law”

$$\frac{d}{dt} \sum_n (z_n P_{nz} + \bar{z}_n P_{n\bar{z}}) = 2(1 - \frac{M}{4\pi}). \quad (30)$$

The fact that the terms depending on g in eq.(25) do not contribute can be easily seen if we write the r.h.s of (25) times z in the form

$$-4z\partial_M Q(z) + \pi^{\bar{z}}_z(z\partial_z g - g) + \partial_z(z\pi^{\bar{z}}_z g). \quad (31)$$

The last term is harmless since it is a total derivative of a rational function; the previous one falls again as $1/z^2$ since the combination $(z\partial_z g - g)$ cancels the linear behavior of g at infinity.

Eq.(28) is a severe constraint on the form of the reduced hamiltonian $\mathcal{H}$ describing the motion of the particles when gravity is eliminated. In fact $\sum_n P_{nz} = 0$ imposes

$$\mathcal{H} = \mathcal{H}(\zeta_2, \dots, \zeta_N, P_2, \dots, P_N) \quad (32)$$

with $\zeta_n = z_n - z_1$ and eq.(28) imposes that

$$\sum_{n>1} \left(P_{nz} \frac{\partial \mathcal{H}}{\partial P_{nz}} - \zeta_n \frac{\partial \mathcal{H}}{\partial \zeta_n} \right) = 1 - \frac{M}{4\pi}. \quad (33)$$

The general solution of this equation has the form

$$\mathcal{H} = \sum_{n>1} \left(\log(P_{nz} P_{n\bar{z}}) + \log(\zeta_n \bar{\zeta}_n)^{\alpha_n} \right) + \mathcal{H}_0 \quad (34)$$

where $\mathcal{H}_0$ is a homogeneous function of degree zero in the variables P_{nz} and $1/\zeta_n$ and $\sum_{n>1} (1 - \alpha_n) = 1 - \frac{M}{4\pi}$. This property will be seen explicitly realized for two particles with $\mathcal{H}_0 = 0$.

8 The two body problem

In the two body problem, $\pi^{\bar{z}}_z$ has no zero i.e. we are in absence of apparent singularities; $g(z) = z/P_{1z}(z_2 - z_1)$ in order to have a non rotating frame at infinity. The simple residues β_1, β_2 are provided by the Fuchs relations in terms of particle masses m_1, m_2 and the total energy. f appearing in eq.(16) is given by the ratio of two hypergeometric functions which provides the metric [6, 7, 14]. On the other hand it is interesting that the particle trajectories can be determined even before the explicit computation of the metric. In fact we have for

the two particles the equation of motion (20,21). The rewriting of the two formulas for the problem at hand gives for $l = z_1 - z_2$ and $P_z = P_{1z} = -P_{2z}$

$$\dot{l} = \frac{1}{P_z}, \quad \dot{P}_z = -\frac{\mu}{l} \quad (35)$$

whose general solution is

$$l = \text{const} [(1 - \mu)(t - t_0) - iL/2]^{\frac{1}{1-\mu}}. \quad (36)$$

The scattering angle is immediately computed from the previous equation to be $\theta_{scatt} = \pi M/(4\pi - M)$ [1, 24]. It is interesting to note that the scattering angle depends only on the total energy and not on the masses of the individual particles. The two equations (35) are generated by the hamiltonian $H_{eff} = \ln(P_x^2 + P_y^2) + \mu \ln(l_x^2 + l_y^2)$ which is unique up to an additive constant. Such an hamiltonian should not be confused with the total energy of the system; it is the effective hamiltonian for the time development of the two particle system and contains as an essential parameter the total energy $M = 4\pi\mu$.

9 The $\mathcal{N}$ body problem

We shall consider in this section the $\mathcal{N}$ -particle case with particular attention to $\mathcal{N} = 3$. In flat space the number of Lorentz invariants is easily seen to be $3\mathcal{N} - 3$ as for $n > 3$ the $2 + 1$ -vector $\mathbf{P}_n$ is determined by the scalar product of $\mathbf{P}_n$ with $\mathbf{P}_1, \mathbf{P}_2, \mathbf{P}_3$. In curved space we have the same number of invariants. As it is well known here the invariants are replaced by the traces of the holonomies around the world lines of an arbitrary collection of particles [1, 25, 26]. The parallel transport monodromy matrices which belong to $SO(2, 1)$ or in the fundamental representation to $SU(1, 1)$ are defined up to a conjugation. Keeping in mind that every $SU(1, 1)$ holonomy has three real degrees of freedom we have $3\mathcal{N}$ degrees of freedom to which we have to subtract the three degrees of freedom of the $SU(1, 1)$ conjugation thus reaching as expected the same number of invariants as in the flat case.

In the present approach parameterizing the dynamics in terms of the position of the particles and their momenta is not a convenient choice, because these, as we have anticipated,

are not the natural variables for the associated Riemann-Hilbert problem. In order to understand the problem better let us consider the case $\mathcal{N} = 3$. It is useful to go over from the equation in projectively canonical form (13) to the equivalent one presented below [23]

$$y'' + py' + qy = 0 \quad (37)$$

with

$$Q = q - \frac{p'}{2} - \frac{p^2}{4}. \quad (38)$$

The transformation can be so chosen as to have the following Riemann scheme

$$\left(\begin{array}{ccccc} 0 & 1 & z_3 & z_A & \infty \\ 0 & 0 & 0 & 0 & \rho_\infty \\ \mu_1 & \mu_2 & \mu_3 & 2 & \rho_\infty + \mu_\infty \end{array} \right) \quad (39)$$

where the Fuchs relation

$$\mu_1 + \mu_2 + \mu_3 + \mu_\infty + 2\rho_\infty = 1 \quad (40)$$

fixes ρ_∞ in terms of the other parameters and

$$p(z) = \frac{1 - \mu_1}{z} + \frac{1 - \mu_2}{z - 1} + \frac{1 - \mu_3}{z - z_3} - \frac{1}{z - z_A};$$

$$q(z) = \frac{\kappa}{z(z - 1)} - \frac{z_3(z_3 - 1)H}{z(z - 1)(z - z_3)} + \frac{z_A(z_A - 1)b_A}{z(z - 1)(z - z_A)}$$

with $\kappa = \rho_\infty(\rho_\infty + \mu_\infty)$. Here we use the unconstrained parameterization of ref.[23]; the connection with the parameters μ_n and β_n can be easily obtained by comparing the residues in eq.(38).

The parameters μ_i and μ_∞ are free real parameters which give the trace of the holonomies around the particles and infinity (total energy). z_A is given in terms of the z_n and the P_n ($n = 1, 2, 3$), while H is given by the no-logarithm condition ⁷ $Q_A(z_A) = \beta_A^2/2$, i.e. the

⁷Here $Q_A(z)$ denotes the function Q without its poles in z_A .

requirement that the point z_A is a regular point for 2σ [23]. Such H is easily found

$$H = \frac{z_A(z_A - 1)(z_A - z_3)}{z_3(z_3 - 1)} \left\{ b_A^2 - \left(\frac{\mu_1}{z_A} + \frac{\mu_2}{z_A - 1} + \frac{\mu_3 - 1}{z_A - z_3} \right) b_A + \frac{\kappa}{z_A(z_A - 1)} \right\}. \quad (41)$$

Thus we have the following free real parameters $\mu_1, \mu_2, \mu_3, \mu_\infty, \text{Re}(z_A), \text{Im}(z_A), \text{Re}(b_A), \text{Im}(b_A)$ which are eight real parameters, to be contrasted with the six real invariants we have enumerated at the beginning of this section. The reason is that the constraint that the holonomies described by eq.(37) belong to $SU(1, 1)$, fixes the value of $\text{Re}(b_A), \text{Im}(b_A)$ in terms of all other parameters. The explicit form of this relation has been investigated in the mathematical literature [28] but we are not aware of any explicit form of it. Adding a new particle introduces a new particle singularity with exponent μ_4 i.e. $p(z)$ acquires a new term of the type $(1 - \mu_4)/(z - z_4)$, but in addition a new apparent singularity is generated say at z_B . The indices of this new singularity have to be integers, 0 and 2 while the new simple residue b_B has to be a function of all other parameters, and in particular of z_B , because the $SU(1, 1)$ character of the holonomies allow only an increase of 3 in the number of free real parameters i.e. $\mu_4, \text{Re}z_B, \text{Im}z_B$. Adding other particles does not alter the procedure.

The complete solution of the dynamical problem requires that we provide the equations of motion for all the variables parameterizing the dynamics of the system and in particular for the apparent singularities and their first order residues. To this purpose, we shall follow here a synthetic approach which will encode in a single equation all the information on the time development of the positions and residues of all singularities. To accomplish this we shall exploit the trace part of equation for $\dot{g}_{ij}$ i.e.

$$\frac{d(2\sigma)}{dt} = N^z \partial_z(2\sigma) + \partial_z N^z + N^{\bar{z}} \partial_{\bar{z}}(2\sigma) + \partial_{\bar{z}} N^{\bar{z}} \quad (42)$$

which combined with the time development of $\pi_{\bar{z}}^z(z)$, eq.(24), provides the time derivative of the reduced conformal factor $\tilde{\sigma}$ i.e.

$$2\dot{\tilde{\sigma}} = g(z) \partial_z(2\tilde{\sigma}) - g'(z) + \text{c.c.} \quad (43)$$

We shall exploit now the fundamental relation (26) of the theory of fuchsian differential equation [23, 21]. Taking the time derivative of eq.(26) we have

$$\dot{Q}(z) = \frac{1}{2}g'''(z) + 2g'(z)Q(z) + g(z)Q'(z). \quad (44)$$

The above equation implies that the time development of $Q(z)$ is given by a conformal diffeomorphism generated by $Q(z)$ itself. We are not in presence of a classical transformation but of an anomalous one. Extensions of the classical Liouville theory have been considered in [27].

It is interesting that this equation, in addition to the particle equations of motion, contains the whole Garnier system for the time evolution of the apparent singularities. We recall that $g(z)$ is the rational function introduced in eq.(21); it is given by a sum of simple poles located at the apparent singularities and of a first order polynomial.

Equating the simple residue of eq.(44) at infinity one simply obtains

$$\frac{d}{dt}(\sum_n \beta_n + \sum_B \beta_B) = 0 \quad (45)$$

which is already contained in the Fuchs relation (23), while equating the second order residues at infinity we obtain

$$\frac{d}{dt} \left[\sum_n (1 - \mu_n^2) + 2 \sum_n \beta_n z_n + 2 \sum_B \beta_B z_B \right] = 0$$

which using the second Fuchs relation (23) provides $d\mu_\infty^2/dt = 0$, where $\mu_\infty = 1 - M/4\pi$ i.e. the conservation of total energy. On the other hand if we equate the second order residues of eq.(44) at the particle pole z_n we obtain $d\mu_n^2/dt = 0$, where $\mu_n = m_n/4\pi$ i.e. the conservation of the individual particle rest masses. Matching of the third order residue on the particle singularity z_n gives

$$\dot{z}_n = -g(z_n) \quad (46)$$

while matching of the first order residue gives $\dot{\beta}_n$ i.e.

$$\dot{\beta}_n = \frac{1 - \mu_n^2}{2} g''(z_n) + g'(z_n) \beta_n. \quad (47)$$

The motion of the apparent singularities is given by the matching of the residues of the third order pole at z_A . If we Laurent expand $g(z)$ around z_A

$$g(z) = \frac{g_{-1}}{z - z_A} + g_0 + g_1(z - z_A) + O((z - z_A)^2) \quad (48)$$

we obtain

$$\dot{z}_A = -g_{-1}\beta_A - g_0 \quad (49)$$

while the first order pole gives

$$\dot{\beta}_A = -2g_{-1}Q'_A(z_A) + g_1\beta_A - \frac{3}{2}g_2. \quad (50)$$

Equations (49) and (50) can be written in hamiltonian form; for example in the three body problem, by choosing $g(z)$ as to cancel the pole in eq.(19) and keep particle 1 fixed at $z_1 = 0$ and particle 2 fixed at $z_2 = 1$, we have

$$\begin{aligned} \frac{\partial z_A}{\partial z_3} &= \frac{z_A(z_A - 1)(z_A - z_3)}{z_3(z_3 - 1)} \times \\ \left[\beta_A - \frac{1}{z_A - 1} - \frac{1}{z_A} \right] &= \frac{\partial H}{\partial b_A} \end{aligned} \quad (51)$$

and

$$\frac{\partial \beta_A}{\partial z_3} = -\frac{\partial H}{\partial z_A}. \quad (52)$$

where H is the so called Garnier hamiltonian and it given by (41). The parameter β_A is connected to those appearing in H by the relation with

$$\frac{\beta_A}{2} = b_A + \frac{1}{2} \left(\frac{1 - \mu_1}{z_A} + \frac{1 - \mu_2}{z_A - 1} + \frac{1 - \mu_3}{z_A - z_3} \right). \quad (53)$$

The explicit expression of H can also be written for the general $\mathcal{N}$ body problem.

It is now possible to interpret the problem of gravity in presence of particles à la Einstein-Infeld- Hoffmann [29] i.e. as pure gravity without particles in which one allows for singular solutions of the field equations. What we found is that the deficit of the singularity has to remain constant (equivalent to the constancy of rest masses), the total angular deficit has to remain constant (equivalent to the constancy of the total energy) and the field singularities follow geodesics of space- time.

10 Conclusions

The application of the instantaneous York gauge $K = 0$, to $2 + 1$ dimensional gravity is practically restricted to the case of open universes; on the other hand in this case the advantages are several and profound. The momenta π^a_b conjugate to the space metric have only two independent components and the diffeomorphism constraint has a very simple solution i.e. the $\pi^{\bar{z}}_z$ ($\pi^z_{\bar{z}}$) are rational functions of z ($\bar{z}$) whose residues are the particle momenta P_n .

The advantage is still greater with regard to the hamiltonian constraint which in this gauge is equivalent to an inhomogeneous Liouville equation for the reduced conformal factor $\tilde{\sigma}$; the sources are the particle singularities i.e. the poles of $\pi^{\bar{z}}_z$ and the apparent singularities, i.e. the zeros of $\pi^{\bar{z}}_z$. The function N is given by the derivative of $\tilde{\sigma}$ with respect to the total energy while N^z is given in terms of N by

$$N^z = -\frac{2}{\pi^{\bar{z}}_z} \partial_z N + g(z), \quad (54)$$

where $g(z)$ is a rational function which cancels the polar singularities of the first term and grows at infinity not faster than z . The linear term in $g(z)$ is fixed if we want to deal with a reference frame which does not rotate at infinity.

The equations of motion for the position of the particles and of the apparent singularities and also the time dependence of the linear residues at such singularities are given by the time development of the function $Q(z)$ entering the fuchsian differential equation which underlies the conformal factor describing the space metric. Such a time development is given by the energy momentum tensor of a conformal Liouville theory and it encodes completely the dynamics of the system.

11 Acknowledgments

We are grateful to Marcello Ciafaloni, Stanley Deser, Marc Henneaux and Roman Jackiw for useful discussions.

References

- [1] S. Deser, R. Jackiw and G. 't Hooft, Ann. Phys. (NY) 152 (1984) 220; S. Deser and R. Jackiw, Ann. Phys. 153 (1984) 405.
- [2] V. Moncrief, J.Math. Phys. 30 (1989) 2907.
- [3] A. Hosoya and K. Nakao, Class. Quantum Grav. 7 (1990) 163.
- [4] G. 't Hooft, Class. Quantum Grav. 9 (1992) 1335; Class. Quantum Grav. 10 (1993) 1023.
- [5] H. Waelbroeck, Class. Quantum Grav. 7 (1990) 751; Nucl. Phys. B364 (1991) 475; Phys. Rev. D50 (1994) 4982.
- [6] A. Bellini, M. Ciafaloni, P. Valtancoli, Physics Lett. B 357 (1995) 532; Nucl. Phys. B 462 (1996) 453.
- [7] M. Welling, Class. Quantum Grav. 13 (1996) 653; Nucl. Phys. B 515 (1998) 436.
- [8] J. Nelson, T. Regge, Phys. Lett. B273 (1991) 213.
- [9] S. Carlip, Phys. Rev. D42 (1990) 2647.
- [10] A. Hosoya and K. Nakao, Progr. Theor. Phys. 84 (1990) 739.
- [11] G. 't Hooft, Class. Quantum Grav. 13 (1996) 1023.
- [12] H. Waelbroeck, Phys. Rev. D50 (1994) 4982.
- [13] M. Welling, Class. Quant. Grav. 14 (1997) 3313; H-J. Matschull and M. Welling, Class. Quantum Grav. 15 (1998) 2981; J. Louko and H-J. Matschull, preprint AEI 1999-015, MZ-TH/99-31, gr-qc/9908025.
- [14] P. Menotti and D. Seminara, Annals of Physics, in press.
- [15] S.W. Hawking and C. J. Hunter, Class. Quantum Grav. 13 (1996) 2735; G. Hayward, Phys. Rev. D 47 (1993) 3275.

- [16] R. Arnowitt, S. Deser and C.W. Misner, in “Gravitation: an introduction to current research” Edited by L.Witten, John Wiley & Sons New York, London 1962.
- [17] R. M. Wald, “General Relativity”, The University of Chicago Press, Chicago and London.
- [18] M. Henneaux, Phys. Rev. D 29 (1984) 2767.
- [19] S. Deser, Class. Quantum Grav. 2 (1985) 489.
- [20] A. Ashtekar and M. Varadarajan Phys.Rev. D50 (1994) 4944.
- [21] L. Liouville J. Math. Pures Appl. 18 (1853) 71; H. Poincaré, J. Math. Pures Appl. 5 ser.2 (1898) 157; P. Ginsparg and G. Moore, hep-th/9304011; L. Takhtajan, “Topics in quantum geometry of Riemann surfaces: two dimensional quantum gravity”, Como Quantum Groups (1994) 541, hep-th/9409088; I. Kra, “Automorphic forms and Kleinian groups” Benjamin, Reading Mass., 1972.
- [22] W.P. Thurston, *The Geometry and Topology of Three-Manifolds*, Princeton lecture notes (1979).
- [23] M. Yoshida, “Fuchsian differential equations”, Fried. Vieweg & Sohn, Braunschweig; K. Okamoto, J. Fac. Sci. Tokio Univ. 33 (1986).
- [24] G. 't Hooft, Comm. Math. Phys. 117 (1988) 685.
- [25] S. M. Carrol, E. Farhi, A. H. Guth and K. O. Olum, Phys. Rev. D 50 (1994) 6190.
- [26] P. Menotti and D. Seminara, Ann. Phys. (N.Y.) 240 (1995) 203.
- [27] T. Curtright and C. Thorn, Phys. Rev. Lett. 48 (1982) 1309; E. D'Hoker, R. Jackiw, Phys. Rev. D26 (1982) 3517.
- [28] L. Schlesinger “Vorlesungen über lineare Differentialgleichungen” and “Handbuch der Theorie der linearen Differentialgleichungen” Teubner Verlag, 1908, Berlin, Leipzig.

- [29] A. Einstein, L. Infeld and B. Hoffmann, *Ann. Math.* 39 (1938) 65; J. N. Goldberg, in “Gravitation: an introduction to current research” Edited by L.Witten, John Wiley & Sons New York, London 1962.